

Cassia grandis

pink shower, carao

L.f.

Fabaceae - Caesalpinioideae

LOCAL NAMES

English (coral shower, apple blossom cassia, pink shower, liquorice tree, horse cassia); French (bâton casse, casse du Brésil); Lao (Sino-Tibetan) (may khoum); Malay (kotek mamak); Spanish (sandal, carao, carámano, cañafistula, cañadonga); Thai (kanpaphruk (Bangkok)); Trade name (pink shower, carao); Vietnamese ([oo] m[oo])

BOTANIC DESCRIPTION

Cassia grandis is a medium-sized tree, up to 20(-30) m tall, semi-deciduous, young branches and inflorescence covered with rusty lanate indumentum.

Leaves with 10-20 pairs of leaflets, petiole 2-3 cm long, lanate, leaflets subsessile, elliptical-oblong, 3-5 cm x 1-2 cm, subcoriaceous, rounded at both ends.

Inflorescence a lateral raceme, 10-20 cm long, 20-40-flowered; flowers with sepals 5-8 mm long, petals initially red, fading to pink and later orange, the median one red with a yellow patch, stamens 10 with hirsute anthers, 3 long ones with filaments up to 30 mm and anthers 2-3 mm long, 5 short ones with filaments 7-9 mm and anthers 1-1.5 mm long, 2 reduced ones with filaments about 2 mm long.

Fruit pendent, compressed, 20-40(-60) cm long, 3-5 cm in diameter, blackish, glabrous, woody, rugose; seeds 20-40 per pod, surrounded by sweetish pulp.

The roots of *C. fistula* and *C. javanica* lack nodulating ability, but for *C. grandis* this is not clear.

BIOLOGY

It is reported evergreen in Java and deciduous in northern Malaysia and Indo-China, where the leaves fall at the beginning of the dry season. The tree flowers before new leaves appear. In Costa Rica, fruit takes 10-12 months to mature.

Cassia grandis

L.f.

Fabaceae - Caesalpinioideae

pink shower, carao

ECOLOGY

C. grandis is a common element of lowland and riparian, semi-deciduous forests.

BIOPHYSICAL LIMITS

Altitude: 0- 600 m

Mean annual rainfall: 1000-2800 mm

Mean annual temperature: 21-26 deg C

Soil type: *C. grandis* tolerates seasonally waterlogged soils.

DOCUMENTED SPECIES DISTRIBUTION

Native: Belize, Bolivia, Brazil, Chile, Colombia, Costa Rica, Cuba, Dominica, El Salvador, Guatemala, Guyana, Haiti, Honduras, Jamaica, Mexico, Nicaragua, Panama, Peru, Puerto Rico, Surinam, Venezuela

Exotic: Cambodia, Cote d'Ivoire, Democratic Republic of Congo, Fiji, India, Indonesia, Laos, Malaysia, Papua New Guinea, Seychelles, Sierra Leone, Sri Lanka, Tanzania, Uganda, Vietnam



The map above shows countries where the species has been planted. It does neither suggest that the species can be planted in every ecological zone within that country, nor that the species can not be planted in other countries than those depicted. Since some tree species are invasive, you need to follow biosafety procedures that apply to your planting site.

Cassia grandis

pink shower, carao

L.f.

Fabaceae - Caesalpinioideae

PRODUCTS

Food: The membrane surrounding seeds is used as a chocolate substitute in Central America.

Fodder: Cattle relish the fruit pods of the tree.

Fuel: The tree is considered good for charcoal and fuelwood.

Timber: *C. grandis* is reported to give strong multipurpose wood, used in joinery, carpentry, beams among others.

Gum or resin: The seeds of *C. grandis* are a potential commercial source of gums. Seed gum is a potential binder for the pharmaceutical industry.

Medicine: The fruit pulp is used as a laxative similar to *C. fistula* and reported to be more powerful. The ripe pods and seeds of *C. grandis* are also used as a laxative. A decoction of the leaves is used as a laxative and in the treatment of lumbago. Fresh juice of the leaves of *C. grandis* is used externally in the treatment of ringworm.

Anthraquinones are found in *C. grandis* (aloe-emodin). Compounds isolated from *C. grandis* include centaureidine, catechin, myristicin, 2,4-dihydroxybenzaldehyde, 3,4,5-trimethoxybenzaldehyde, 2,4,6-trimethoxybenzaldehyde, beta-sitosterol, kokusaginine (6,7-dimethoxyfuroquinoline) and fabioline (1,1'-bipiperidine). The ethanol extract of the leaves and bark of *C. grandis* showed in vitro antifungal activity against *Epidermophyton floccosum*, *Microsporium gypseum* and *Trichophyton rubrum* in pure culture at a minimal inhibitory concentration of 50 microgram/ml.

SERVICES

Reclamation: The tree is recommended for revegetation in especially periodically flooded areas.

Ornamental: *C. grandis* has been widely introduced for ornamental purposes.

Boundary or barrier or support: *C. grandis* can be planted as a live fence.

Intercropping: *C. grandis* is recommended for dry zone intercropping with perennial crops and in pastures.

Cassia grandis

pink shower, carao

L.f.

Fabaceae - Caesalpinioideae

TREE MANAGEMENT

C. grandis requires regular pruning. It has the ability to sucker.

GERMPLASM MANAGEMENT

Seed can be stored for prolonged periods without loss of viability.

PESTS AND DISEASES

Cassia grandis

pink shower, carao

L.f.

Fabaceae - Caesalpinioideae

FURTHER READING

CABI. 2000. Global Forestry Compendium. CD-ROM. CABI

Farooqi MIH, Kapoor VP & Islam G. 1978. Seeds of genus *Cassia* as possible sources of industrial gums. *Indian Forester*. 104(11): 729-733.

Flores EM, Rivera DI & Vasquez NM. 1986. Germinacion y desarrollo de la plantula de *Cassia grandis* L. (Caesalpinioideae) [Germination and development of the seedling of *Cassia grandis* L. (Caesalpinioideae)]. *Revista de Biologia Tropical*. 34(2): 289-296.

Irwin HS & Barneby RC. 1982. The American Cassiinae. A synoptical revision of Leguminosae tribe Cassieae subtribe Cassiinae in the New World. *Memoirs of the New York Botanical Garden*. 35(2): 64-635.

Jolin D & Torquebiau E. 1992. Large cuttings: a jump start for tree planting. *Agroforestry Today*. 4(4): 15-16.

Nguyen Van Duong. 1993. Medicinal plants of Vietnam, Cambodia and Laos. Mekong Printing, Santa Ana, California, United States. 528pp.

Rocas AN. 2002 *Cassia grandis* L.f. In: Vozzo, J.A. (ed.) Tropical tree seed manual. U.S. Department of Agriculture, Forest Service. Agriculture Handbook No. 721. Washington, USA. pp. 369-371.

Valencia E, Madinaveitia A, Bermejo J, Gonzalez AG & Gupta MP. 1995. Alkaloids from *Cassia grandis*. *Fitoterapia*. 66(5): 476-477.

SUGGESTED CITATION

Orwa C, A Mutua, Kindt R, Jamnadass R, S Anthony. 2009 *Agroforestry Database: a tree reference and selection guide version 4.0* (<http://www.worldagroforestry.org/sites/treedbs/treedatabases.asp>)