

Cassipourea malosana

ndiri, muzaizi, msengera

Alston

Rhizophoraceae

LOCAL NAMES

English (pillarwood); Trade name (ndiri,muzaizi,msengera)

BOTANIC DESCRIPTION

Cassipourea malosana is an evergreen tree approximately 6-45 m in height. It bears a conspicuously straight, pale, cylindrical trunk and a small rounded crown. The branches radiate out at right angles to the trunk. Bark smooth, pale grey with well defined ring marks. Slash soft, pale orange with a darker outer edge.

Leaves opposite, dark green, 5 x 3 cm, but variable in size; glossy, tough and borne on woody twigs, margins toothed or entire.

Flowers 1-5(-8) per inflorescence, laterally placed under leaves; pedicels up to 4 mm, rarely 7 mm long, densely puberulous. Calyx 4-5-partite; tube 1 mm long; lobes 3.5-5 mm long, narrowly oblong-triangular, apex acute, densely appressed-pubescent externally, glabrous or very nearly so within. Petals 4-5 lacinate apically, anthers introrse. Ovary superior, 3-4 celled, glabrous or almost so with a few apical hairs.

Fruit a thin fleshy dehiscent capsule, 1 cm long, becoming black when dry.

The species is quite variable with regard to leaf morphology. The specific epithet malosana is after Mt. Malosa in Malawi where the type specimen was collected.

Cassipourea malosana

ndiri, muzaizi, msengera

Alston

Rhizophoraceae

ECOLOGY

C. malosana is a tree commonly found in moist montane forest in association with *Podocarpus* spp. It also occurs in upland dry forest.

BIOPHYSICAL LIMITS

Altitude: 750-2 550 m

Mean annual rainfall: over 2 200 mm

DOCUMENTED SPECIES DISTRIBUTION

Native: Ethiopia, Kenya, Malawi, Mozambique, Somalia, Sudan, Tanzania, Uganda

Exotic:



The map above shows countries where the species has been planted. It does neither suggest that the species can be planted in every ecological zone within that country, nor that the species can not be planted in other countries than those depicted. Since some tree species are invasive, you need to follow biosafety procedures that apply to your planting site.

Cassipourea malosana

ndiri, muzaizi, msengera

Alston

Rhizophoraceae

PRODUCTS

Fuel: Tree parts used as fuelwood.

Timber: *C. malosana* is an important timber tree in East Africa with undifferentiated sapwood and heartwood, white to light brown often with purplish streaks resulting from fungal attack. Specific gravity 0.59. Texture fine and even; grain usually straight but with a slight to marked tendency to spiraling. It has very hard timber which dries slowly and is subject to distortion. Timber suited for flooring, turnery, tool handles and construction work requiring great strength and elasticity. Sawing of green timber is difficult because of its tendency to spring. Dry wood is easier to work.

SERVICES

Erosion control: The tree is found in catchment areas (wet montane forest) and contributes in soil conservation of these areas.

Shade or shelter: *C. malosana* provides shade.

Soil improver: Experimental results indicate Potassium is more easily leached from *C. malosana* leaves than calcium or magnesium.

Cassipourea malosana

ndiri, muzaizi, msengera

Alston

Rhizophoraceae

Cassipourea malosana

ndiri, muzaizi, msengera

Alston

Rhizophoraceae

FURTHER READING

Beentje HJ. 1994. Kenya trees, shrubs and lianas. National Museums of Kenya.

<http://www2.fpl.fs.fed.us/TechSheets/Chudnoff/African/htmlDo.../Cassipoureamalosan>

Lewis J. 1956. Rhizophoraceae. In: Flora of Tropical East Africa. Crown Agents, London.

Sommerlatte H. & Sommerlatte M. 1990. A field guide of the trees and shrubs of the Imatong Mountains of Southern Sudan. Deutsche Gesellschaft für Technische Zusammenarbeit, Nairobi.

Zamierowski EE. 1975. Leaching losses of minerals from leaves of trees in montane forest in Kenya. *Journal of Ecology*. 63(2): 679-687.

SUGGESTED CITATION

Orwa C, A Mutua, Kindt R, Jamnadass R, S Anthony. 2009 *Agroforestry Database: a tree reference and selection guide version 4.0* (<http://www.worldagroforestry.org/sites/treedbs/treedatabases.asp>)