

### LOCAL NAMES

English (magic gwarra); Luganda (nsikizi); Swahili (mdaa); Tswana (motlhakola); Zulu (umhlangula)

### BOTANIC DESCRIPTION

*Euclea divinorum* is a shrub or small tree up to about 6 m tall, often branching from the base or sometimes with a single stem. Bark grey, fairly smooth in young trees but fissured in older specimens. Crown much branched and grey-green in colour.

Leaves simple, coriaceous, lanceolate, margins wavy, sub-opposite or alternate, 3.5-9 cm long and 1-2.5 cm wide. Upper surfaces light green or grey green, sometimes with a yellowish tinge, lower surface pale and smoother in texture. Nerves visible as fine lines and midrib raised below.

Flowers small, cup-shaped and creamy in colour borne on a short dense head, flowers and inflorescence covered with tiny, rusty-brown dots. Male and female flowers on separate trees.

Fruit a round, thinly fleshed berry, usually 1-seeded, purple when ripe.

The botanical author of the species, William Phillip Hiern noted the popular use of the plant by diviners and thus coined the specific epithet *divinorum* for it.

### BIOLOGY

*E. divinorum* is dioecious. The fruits are at times dispersed by Hornbills.



*E. divinorum* leaves (Bob Bailis)



*E. divinorum* bark (Bob Bailis)



Young *E. divinorum* (Bob Bailis)

**ECOLOGY**

*E. divinorum* is a species common in bush, dry forest margins, thornscrub and open woodlands. It is usually associated with *Acacia* spp. and also grows on anthills and river banks in hot dry areas below 900 m. Inter-specific competition between *E. divinorum* and *A. nilotica* leads to conversion of open grasslands to densely vegetated woodlands. The successional dynamics of *Acacia-Euclea* savannahs appear to depend on the browsing of *A. nilotica* and other *Acacia* spp. by large mammals especially elephants.

**BIOPHYSICAL LIMITS**

Altitude: 1-2 400 m

Mean annual temperature: 17 deg C

Mean annual rainfall: 700 mm

Soil type: Prefers mesic calcareous valley clays and sometimes rocky ground.

**DOCUMENTED SPECIES DISTRIBUTION**

Native: Botswana, Kenya, Namibia, South Africa, Swaziland, Tanzania, Uganda, Zimbabwe

Exotic:



The map above shows countries where the species has been planted. It does neither suggest that the species can be planted in every ecological zone within that country, nor that the species can not be planted in other countries than those depicted. Since some tree species are invasive, you need to follow biosafety procedures that apply to your planting site.

**PRODUCTS**

**Food:** The fruits are edible. In East Africa the bark is used in the preparation of fatty-meat and milk soups. The roots are chewed to impart a red colour to the mouth.

**Fodder:** The leaves are browsed by wild animals such as the rhino, giraffe and grey duiker.

**Fuel:** Branches used as firewood in some East African localities.

**Tannin or dyestuff:** The bark is rich in phenolics (123 mg/g) and tannins(94 mg/g). The bark extract is used in dyeing baskets woven from *Hyphaene petersiana* (mokola palm) leaves. Leather produced from *E. divinorum* has a deep red colour that provides an aesthetically pleasing alternative to black wattle and other available barks. The root is used in the production of black floor mats. In Ovamboland a purple ink is made by boiling the fruit.

**Alcohol:** The fruit is used in making beer.

**Poison:** Tannic and gallic acid extracts in this plant may contain sufficient amounts of inhibitory constituents to interfere with the virulence and growth of enamel cavity causing bacteria *in vivo*.

**Medicine:** *E. divinorum* branches are used as chewing sticks for oral care. The fruits are said to be strongly purgative. In Kenya the root decoction is used as a purgative and the bark infusion as an appetizer. Decoctions of the root are used by the Zulu for toothache.

**Other products:** The pentacyclic triterpenes and naphthoquinones, lupeol, lupene, betulin, 7-methyljuglone, isodiospyrin, shinalone, catechin and the novel 3 beta-(5-hydroxyferuloyl) lup-20(30)-ene were isolated from roots of *E. divinorum* collected in Zimbabwe.

**SERVICES**

**Erosion control:** The species is important in protecting soil.

**Shade or shelter:** The plant is a good shade tree.

**Reclamation:** Because of its tolerance of high arsenic soil levels *E. divinorum* can be used in reclamation of gold mine pit wastes.

**Other services:**

In South Africa the tree is used to repel witches and other evil, the branches are hung above entrances into huts and kraals.

**TREE MANAGEMENT**

The plant has remarkable coppicing and root suckering ability and if not checked, tends to weediness, dominating pasture to the detriment of wildlife and pastoralism. Use of arboricides is recommended in weedy situations.

**FURTHER READNG**

Ben-Shahar R. 1991. Successional patterns of woody plants in catchment areas in a semi-arid region. *Vegetatio*. 93(1): 19-27.

Cunningham AB and Milton SJ. 1987. Effects of basket-weaving industry on mokola palm and dye plants in northwestern Botswana. *Economic Botany*. 41(3): 386-402.

Homer KA, Manji F and Beighton D. 1990. Inhibition of protease activities of periodontopathic bacteria by extracts of plants used in Kenya as chewing sticks (mswaki). *Archives of Oral Biology*. 35(6): 421-424.

Mebe PP, Cordell GA and Pezzuto JM. 1998. Pentacyclic triterpenes and naphthoquinones from *Euclea divinorum*. *Phytochemistry*. 47(2): 311-313.

Palmer E, Pitman N. 1972. *Trees of Southern Africa Vol. 2*. A.A. Balkema Cape Town.

van Grinsven M, Parkipuny ML and Johns T. 1999. *Euclea divinorum* (Ebenaceae) bark is a high-potential tanning material. *Economic Botany*. 53: 220-221.

Wild H. 1974. Geobotanical anomalies in Rhodesia 4. The vegetation of arsenical soils. *Kirkia*. 9(2): 243-264.

**SUGGESTED CITATION**

Orwa C, A Mutua, Kindt R, Jamnadass R, S Anthony. 2009 *Agroforestry Database: a tree reference and selection guide version 4.0* (<http://www.worldagroforestry.org/sites/treedbs/treedatabases.asp>)