

**LOCAL NAMES**

English (false coffee tree); Filipino (kukodmon (Bikol)); Indonesian (taji ayam, melingu (Javanese), engkudug biang); Malay (setebal (Peninsular)); Thai (phawa nam)

**BOTANIC DESCRIPTION**

*Fagraea racemosa* is a shrub or small to medium-sized tree up to 25(-40) m tall, bole up to 30 cm in diameter, without buttresses, bark surface smooth but becoming narrowly and deeply fissured, pale grey to dark grey-brown, inner bark yellow-brown.

Leaves very variable, from broadly ovate via elliptical to obovate-oblong, lanceolate or rarely even linear, 5-50 cm x 1-23 cm, apex rounded to acute and often short-to long- acuminate, secondary veins distinctly prominent below, petiole 0.2-5 cm long, stipules connate into an ocrea surrounding the stem.

Inflorescence terminal and usually drooping, pedicel with bracteoles at the base; corolla tube funnel-shaped, 2-4 cm long, stigma faintly 2-lobed.

Fruit subglobose to ellipsoid-ovoid, apiculate, bluish or greenish or red when ripe.

*F. racemosa* is highly variable and several forms have been distinguished. It develops according to the architectural tree model of Roux, having a monopodial orthotropic trunk with continuous growth and plagiotropic branches, but never by apposition.

**BIOLOGY**

Flowering is sporadic with the main crop in Oct.-Dec.

**ECOLOGY**

*F. racemosa* is found in light to dense primary but more often secondary forest on swampy to dry soil, along rivers but also on podzolized sands, in savannas and lalang grassland vegetation. Locally, it is a conspicuous element of early secondary forest on wastelands and poor soils.

**BIOPHYSICAL LIMITS**

Altitudes: up to 100 m

**DOCUMENTED SPECIES DISTRIBUTION**

Native: Australia, Cambodia, India, Indonesia, Malaysia, Myanmar, Papua New Guinea, Solomon Islands, Vietnam

Exotic:



The map above shows countries where the species has been planted. It does neither suggest that the species can be planted in every ecological zone within that country, nor that the species can not be planted in other countries than those depicted. Since some tree species are invasive, you need to follow biosafety procedures that apply to your planting site.

**PRODUCTS**

Fuel: It is used as firewood.

Timber: The density of the wood is 700-870 kg/m cubic at 15% moisture content. The wood is used as tembesu, but the timber is often only available in small dimensions. It is used for general construction and combs.

Medicine: A decoction of leaves, bark, roots and flowers is used for medicinal purposes, mainly as a tonic after fever and for pains in the loins, but in the Philippines also as an antidote against snake bites. Wood-tar is used to blacken teeth.

Other products: In Papua New Guinea, leaves are used for sealing stone ovens and for wrapping food.

**SERVICES**

Boundary or barrier or support: *F. racemosa* is planted as a live fence.

**TREE MANAGEMENT**

Husbandry: Double stems should be removed. Thinning in cycles of 5 years up to the age of 30 years is appropriate for plantations; thereafter every 10 years. The lower branches are very persistent and pruning these promotes height growth. *F. racemosa* reproduce easily by means of root suckers and coppice freely; locally, trees may be often pollarded for poles.

**GERMPLASM MANAGEMENT**

**PESTS AND DISEASES**

**FURTHER READNG**

Ashton PS. 1988. Manual of the non-dipterocarp trees of Sarawak. Vol. 2. Dewan Bahasa dan Pustaka, Kuala Lumpur. 490 pp.

Browne FG. 1955. Forest trees of Sarawaka and Brunei and their products. Government Printing Office, Kuching. 369 pp.

Cockburn PF. 1976-1980. Trees of Sabah. 2 volumes. Sabah Forest Records No 10. Forest Department Sabah, Sandakan.

Conn BJ & Brown EA. 1996. In Orchard, A.E. (Ed), Flora of Australia 28: 69.

CSIR. 1972. The Wealth of India: A dictionary of Indian raw materials and industrial products, Vol. IX. CSIR.

Henderson CP and Hancock IR. 1988. A guide to the useful plants of Solomon Islands, Research Dept., Ministry of Agriculture and Lands, Honiara, Solomon Islands.

Hildebrand JW et al. 1995. Fagraea Thunb. In Lemmens, R.H.M.J., Soerianegara, I. & Wong, W.C. (Eds.): Plant Resources of South-East Asia No 5(2). Timber trees: Minor commercial timbers. Prosea Foundation, Bogor, Indonesia. pp 234-242.

**SUGGESTED CITATION**

Orwa C, A Mutua, Kindt R, Jamnadass R, S Anthony. 2009 Agroforestry Database: a tree reference and selection guide version 4.0 (<http://www.worldagroforestry.org/sites/treedbs/treedatabases.asp>)