

urucurana, suradan pilon, pilón

**LOCAL NAMES**

English (bully tree,suradan pilon); Portuguese (urucurana mirim,urucurana de leite,aricurqua,sangue-de-boi,licurana,mará-goncalo,marconcal,aricurana,muiragonçalo); Spanish (pilón,cargamanto,casaco,nancitón,pantano,zapatero,sangre de toro,trompillo,palo curtidor); Trade name (pilón,suradan pilon,urucurana)

**BOTANIC DESCRIPTION**

*Hyeronima alchorneoides* is a tall dioecious tree of lowland tropical forests up to 50 m tall and diameter of 1-1.2 m with multiple, orthotropic branches that form a dense crown. The bole tend to be straight, robust, with very little taper, and un-branched up to a height of 20 m or more. Bark smooth, distinctive deep reddish-brown on the outside, pinkish on the inside and bitter-tasting.

Leaves simple, alternate, petiolate, entire, 5-9 cm long, very variable in shape, with large, leafy stipules and abundant peltate hairs on both surfaces. Young leaves green, later turning yellow and nearly red before shedding.

Flowers small and inconspicuous, yellow-green, zygomorphic, arranged on 5 cm long axillary-branched panicles

Fruit a small, one-seeded drupe, 2.5-5.5 mm in diameter, red or dark purple at maturity

Seed very small, surrounded by the stony endocarp, endosperm oily and large embryo.

*H. alchorneoides* is one of 36 species in the genus *Hyeronima* (also known as *Hieronyma*), all of which are indigenous to Central and South America. The species exhibits a wide range of intra-specific variation. Two of the characters most commonly used to identify the tree include leaf size and the presence or absence of stipules. This varies among individuals and even within the same individual, depending on shoot phenology and age.

**BIOLOGY**

*H. alchorneoides* is evergreen with leaves emerging and abscising continuously. The tree is readily recognized in the forest due to the presence of red, pre-senescent leaves in the canopy all year round.

Trees are reproductively mature by the time they are 30 cm in dbh. In many places, flowering and fruiting takes place twice a year. The time of flowering can vary according to rainfall and altitude and sometimes there seems to be no fixed seasonality of the seed production.

In Costa Rica, the trees flower in May-July and sporadically in November-January. The peak of seed production is in January-March. Small insects pollinate the flowers and the species has obligate cross-pollination. The fruits are fleshy, sweet mesocarp, and are dispersed by monkeys and birds

## Hyeronima alchorneoides

Allemão

Euphorbiaceae

urucurana, suradan pilon, pilón

### ECOLOGY

H. alchorneoides is a massive canopy emergent tree of lowland humid and very humid mixed tropical forests, which are sometimes waterlogged during the rainy season.

### BIOPHYSICAL LIMITS

Altitude: 0-900 m

Mean annual temperature: 20-30°C

Mean annual rainfall: 2000-6000 mm

Soil type: H. alchorneoides occurs naturally on alluvial, clayey or acid soils but in plantations, it has successfully grown on older, more weathered ultisols.

### DOCUMENTED SPECIES DISTRIBUTION

Native: Belize, Bolivia, Brazil, Colombia, Costa Rica, Cuba, Ecuador, El Salvador, Guatemala, Guyana, Honduras, Mexico, Nicaragua, Panama, Peru, St Lucia, Surinam, Trinidad and Tobago, Venezuela

Exotic:



The map above shows countries where the species has been planted. It does neither suggest that the species can be planted in every ecological zone within that country, nor that the species can not be planted in other countries than those depicted. Since some tree species are invasive, you need to follow biosafety procedures that apply to your planting site.

**PRODUCTS**

Timber: *H. alchorneoides* has a reddish-brown heartwood, and sapwood with a lighter colour and pinkish hue. The wood specific gravity ranges from 0.59-0.86. It is suitable for heavy construction, including railway sleepers, bridges, columns and beams, fence posts, stakes, piles and in marine construction. In addition, it is used for furniture, cabinetwork, flooring, decorative veneers, turnery and joinery. It is durable and fairly resistant to termites but difficult to impregnate and prone to decomposition, especially below ground

Medicine: The seeds of *H. alchorneoides* contain oil that is effective against intestinal parasites (anti-helminthic). In Guyana, a decoction of the bark is used for its anti-tussive properties.

**SERVICES**

Reclamation: *H. alchorneoides* is among several species that have shown potential for growth on a variety of soils, indicating its suitability for use in re-vegetation of degraded lands. This species has a relatively high allocation to fine roots, making it less likely to experience nutrient limitation under low-fertility conditions. In addition to its tolerance to infertile soils, it grows on acid soils and has a high tolerance for exchangeable aluminium.

Soil improver: *H. alchorneoides* has a role in soil amelioration that may lead to an increase in soil organic carbon from to its large litter production

**TREE MANAGEMENT**

Site preparation prior to stand establishment usually involves site clearing. Potted seedlings transplanted with their nursery substrate and fertilizer applied to the bottom of the planting hole perform better. In Costa Rica, a spacing of 3 x 3 m is used to establish a plantation but 2 x 2 m can also be used. Fertilizer application and weeding in the first 12 months after planting out from the nursery have significant positive effects. First pruning is recommended at 9-12 months while thinning after 3-4 years. Thinned stumps are known to re-sprout, but the resulting root systems tend to be weak.

Trees show good growth and form, and have ability to self-prune. A high shade tolerance by lower leaves make the species compatible for planting in mixed species plantations, where *H. alchorneoides* forms the lower stratum in a stratified canopy. In monocultures, *H. alchorneoides* achieves 90% interception of incoming solar radiation at 2 years of age suppressing understorey regeneration.

**GERMPLASM MANAGEMENT**

It is best to collect the fruits from the tree by cutting down branches. The fruits are small and therefore it is useful to put tarpaulins on the ground under the tree. The fruits are collected when they have changed from green to red, indicating maturity. Once mature, the fruits start to fall and often there are only 3-4 days to carry out the collection

The fruits can be de-pulped although this is not done in many places. The seed storage behaviour is intermediate. The seeds are tolerant to desiccation and low temperatures, and should be stored at low moisture content in airtight bags in a cold. There are about 26000 seeds/kg.

**PESTS AND DISEASES**

The majority of the pests identified for *H. alchorneoides* are defoliating insects. These include *Atta cephalotes*, *Eacles imperialis*, *Exophthalmus*, *Hylesia alinda*, *Megalopyge albicollis*, *Oiketicus abotti*, *Phobetron hipparchia*, *Rothschildia lebeau*, *Spodoptera* spp., *Taeniopoda* and *Trigona* spp.

In addition to defoliators, *H. alchorneoides* is very susceptible to damping off due to fungal attack (*Phytophthora* spp., *Fusarium* spp.) at the seedling stage under very humid nursery conditions.

**FURTHER READING**

- Butterfield RP, Espinoza C M. 1995. Screening trial of 14 tropical hardwoods with an emphasis on species native to Costa Rica: fourth year results. *New Forests*. 9(2): 135-145.
- Butterfield RP. 1995. Promoting biodiversity: advances in evaluating native species for reforestation. *Forest Ecology and Management*. 75(1-3): 111-121.
- Butterfield RP. 1996. Early species selection for tropical reforestation: a consideration of stability. *Forest Ecology and Management*. 81(1-3): 161-168.
- Clark DB, Clark DA, Read JM. 1998. Edaphic variation and the mesoscale distribution of tree species in a neotropical rain forest. *Journal of Ecology*. 86(1): 101-112.
- Clark DB, Clark DA. 1992. Life history diversity of canopy and emergent trees in a neotropical rain forest. *Ecological Monographs*. 62: 315-344.
- Clark DB, Clark DA. 1996. Abundance, growth and mortality of very large trees in neotropical lowland rain forest. *Forest Ecology and Management*. 80(1-3): 235-244.
- Fisher RF. 1995. Amelioration of degraded rain forest soils by plantations of native trees. *Soil Science Society of America Journal*. 59(2): 544-549.
- Franco P. 1990. The genus *Hyeronima* (Euphorbiaceae) in South America. *Botanische Jahrbücher*. 111: 297-346.
- Gerwing JJ. 1995. Competitive effects of three tropical tree species on two species of *Piper*. *Biotropica*. 27(1): 47-56.
- González E. 1992. Humedad y germinación de semillas de *Hyeronima alchorneoides* (Euphorbiaceae). *Revista de Biología Tropical*. 40: 139-141.
- Gonzalez W, Paniagua A, Hernandez Y, Rodriguez JA, Montes de Oca P. 1996. Response of pylon tree (*Hyeronia alchorneoides*) during the nursery stage to the application of three sources of organic matter. *Agronomia Costarricense*. 20(1): 47-51.
- Haggar JP, Ewel JJ. 1995. Establishment, resource acquisition, and early productivity as determined by biomass allocation patterns of three tropical tree species. *Forest Science*. 41(4): 689-708.
- Russo RO, Sandi CL. 1995. Early growth of eight native timber species in the humid tropic region of Costa Rica. *Journal of Sustainable Forestry*. 3(1): 81-84.
- Tinto WF, Blyden G, Reynolds WF, McLean S. 1991. Constituents of *Hyeronima alchorneoides*. *Journal of Natural Products*. 54 (5): 1309-1313.

**SUGGESTED CITATION**

Orwa C, A Mutua, Kindt R, Jamnadass R, S Anthony. 2009 *Agroforestry Database: a tree reference and selection guide version 4.0* (<http://www.worldagroforestry.org/sites/treedbs/treedatabases.asp>)