

Ziziphus mauritiana

jujube

LOCAL NAMES

Amharic (kurkura); Arabic (nabak (fruit),sidr); Bengali (ber boroi,kool,ber,boroi); Burmese (zee-pen,zizidaw,eng-si); English (dunks,jujube,Indian cherry,Indian jujube,Indian plum,geb,ber,common jujube,Chinese date,Chinese apple,bear tree,desert apple); Filipino (manzanita); French (jujube,jujubier,jujubier commun,le jujubier,le jujubier sauvage,liane croc-chien); German (Indischer Jujubenstrauch); Gujarati (bordi); Hindi (baer,badari,elladu,ber,khati,jelachi); Indonesian (widara,dara,bidara); Khmer (putrea); Lao (Sino-Tibetan) (than); Malay (bidara,jujub,epal siam); Mandinka (tomborongo,tomboron moussana,toboro); Nepali (bayer); Sanskrit (kuvala,karkandhu,badara,ajapriya,madhuraphala); Somali (geb,gub); Spanish (yuyuba,Ponseré,perita haitiana); Swahili (mkunazi); Tamil (elandai,yellande); Thai (ma thong,ma tan,phutsan); Tigrigna (geva); Trade name (jujube); Urdu (ber); Vietnamese (tao nhuc,tao,c[aa]y t[as]o ta)

BOTANIC DESCRIPTION

Ziziphus mauritiana is a spiny, evergreen shrub or small tree up to 15 m high, with trunk 40 cm or more in diameter; spreading crown; stipular spines and many drooping branches. Bark dark grey or dull black, irregularly fissured. Where climatic conditions are severe, it is commonly a compact shrub only 3-4 m tall.

Leaves variable, alternate, in 2 rows, oblong-elliptic, 2.5-6 x 1.5-5 cm, with tip rounded or slightly notched base; finely wavy-toothed on edges, shiny green and hairless above; dense, whitish, soft hairs underneath.

Inflorescence axillary cymes, 1-2 cm long, with 7-20 flowers; peduncles 2-3 mm long; flowers 2-3 mm across, greenish-yellow, faintly fragrant; pedicels 3-8 mm long; calyx with 5 deltoid lobes, hairy outside, glabrous within; petals 5, subspathulate, concave, reflexed.

Fruit a drupe, globose to ovoid, up to 6 x 4 cm in cultivation, usually much smaller when wild; skin smooth or rough, glossy, thin but tough, yellowish to reddish or blackish; flesh white, crisp, juicy, subacid to sweet, becoming mealy in fully ripe fruits. Seed a tuberculate and irregularly furrowed stone, containing 1-2 elliptic brown kernels each 6 mm long.

The name 'Ziziphus' is often erroneously written as Zizyphus. The generic name is derived from the latinized version of the Arabic vernacular name 'zizouf' for *Z. jujuba*.

BIOLOGY

Some cultivars attain anthesis early in the morning, others do so later in the day. The flowers are protandrous. Hence, fruit set depends on cross-pollination by insects attracted by the fragrance and nectar. The pollen of the flower is described as 'heavy and thick'. In India, different species of honeybees (*Apis* spp.) and house flies (*Musca domestica*) are reported to be important pollinators; the wasps *Polistes hebraeus* and *Physiphora* spp. have also been observed on flowers. Cross-incompatibility occurs, and cultivars have to be matched for good fruit set; some cultivars produce good crops parthenocarpically. Fruit development takes 4 months in early cultivars to 6 months in late ones. In Southeast Asia, *Z. mauritiana* flowers concurrently with shoot growth in the wet season. Mammals and birds disperse the fruits.

Lam.

Rhamnaceae



Z. mauritiana fruit and leaves. (Anthony Simons)



Z. mauritiana cuttings, Makoka, Zambia (Hannah Jaenicke)



Ziziphus mauritiana slash (Joris de Wolf, Patrick Van Damme, Diego Van Meersschaut)

ECOLOGY

Z. mauritiana is a hardy tree that copes with extreme temperatures and thrives under rather dry conditions. Fruit quality is best under hot, sunny and dry conditions, but there should be a rainy season to support extension growth and flowering, ideally leaving enough residual soil moisture to carry the fruit to maturity. Commercial cultivation usually extends up to 1000 m. Beyond this elevation trees do not perform well, and cultivation becomes less economical. Native to the tropical and subtropical regions, *Z. mauritiana* is more widespread in areas with an annual rainfall of 300-500 mm. It is known for its ability to withstand adverse conditions, such as salinity, drought and waterlogging.

BIOPHYSICAL LIMITS


Altitude: 0-1 500 m, Mean annual temperature: 7-13 to 37-48 deg. C, Mean annual rainfall: 120-2 200 mm

Soil type: Best soils are sandy loam which may be neutral or even slightly alkaline. Can grow on a variety of soils including laterite, black cotton and oolitic limestone.

DOCUMENTED SPECIES DISTRIBUTION

Native: Afghanistan, Algeria, Australia, Bangladesh, China, Egypt, India, Indonesia, Iran, Kenya, Libyan Arab Jamahiriya, Malaysia, Nepal, Pakistan, Thailand, Tunisia, Uganda, Vietnam

Exotic: Angola, Barbados, Botswana, Burkina Faso, Cambodia, Cameroon, Chad, Congo, Cote d'Ivoire, Ethiopia, Gambia, Ghana, Grenada, Guinea, Lesotho, Madagascar, Malawi, Mali, Mauritania, Mozambique, Namibia, Niger, Nigeria, Philippines, Senegal, Sierra Leone, Somalia, South Africa, Sudan, Swaziland, Tanzania, Zambia, Zimbabwe



The map above shows countries where the species has been planted. It does neither suggest that the species can be planted in every ecological zone within that country, nor that the species can not be planted in other countries than those depicted. Since some tree species are invasive, you need to follow biosafety procedures that apply to your planting site.

PRODUCTS

Food: Fruit is eaten fresh or dried and can be made into a floury meal, butter, or a cheeselike paste, used as a condiment. Also used for candy making and pickling. The fruit is a good source of carotene, vitamins A and C, and fatty oils. A refreshing drink is prepared by macerating fruits in water. In Indonesia, young leaves are cooked as a vegetable.

Fodder: In parts of India and North Africa, the leaves of *Z. mauritiana* are used as nutritious fodder for sheep and goats. Analysis of the chemicals constituents on a dry weight basis indicates the leaves contain 15.4% crude protein, 15.8% crude fibre, 6.7% total minerals, and 16.8% starch. In India, the leaves are also gathered to feed tasar silkworms; tasar silk, highly prized, is the only silk commercially exploited in the tropics.

Fuel: *Z. mauritiana* produces excellent firewood (sapwood has 4900 kcal/kg) and good charcoal. Its drooping branches are easily accessible for harvesting.

Apiculture: When in bloom it is occasionally a source of pollen, at best a minor one.

Timber: *Z. mauritiana* yields a medium-weight to heavy hardwood with a density of 535-1080 kg/m³. Heartwood is buff-coloured, pale red or brown to dark brown, sometimes banded or with dark streaks, not sharply demarcated from pale brown sapwood; grain straight, occasionally wavy; texture fine to coarse; wood fairly lustrous. It seasons well but may split slightly during seasoning; easy to work and takes a high finish. It is hard and strong. The wood is used for general construction, furniture and cabinet work, tool handles, agricultural implements, tent pegs, golf clubs, gun stocks, sandals, yokes, harrows, toys, turnery, household utensils, bowling pins, baseball bats, chisels and packaging. It is also suitable for the production of veneer and plywood. Basically, any product that needs a durable, close-grained wood can be made from it.

Tannin or dyestuff: The bark, including the root bark, has served in tanning; when pounded and mashed in water, it yields brown and grey or reddish dyes.

Alcohol: A raw, intoxicating spirit is occasionally distilled from the fermented fruit pulp.

Poison: *Z. mauritiana* is used to stupefy fish in Ethiopia.

Medicine: Leaves, fruits and bark are used medicinally. Pounded roots are added to drinking water and given to poultry suffering from diarrhoea and to humans for indigestion.

Other products: In India, *Z. mauritiana* trees are a host for the lac insects, *Kerria lacca*, which are found on the leaves and make an orange-red resinous substance. The purified resin makes the high-quality ber shellac that is used in fine lacquer work and to produce sealing wax and varnish.

SERVICES

Erosion control: A suitable species to aid in fixation of coastal dune sand.

Shade or shelter: The tree has been planted for shade and windbreaks.

Reclamation: Can withstand severe heat, frost and drought; hence it is planted in dry areas and on sites unfit for other crops.

Ornamental: *Z. mauritiana* is well suited for homegardens.

Boundary or barrier or support: Tree useful as a living fence; its spiny stems and branches deter livestock.

TREE MANAGEMENT

Z. mauritiana is a fast-growing species. Under favourable conditions, height increment on loose soil is 75 cm in 1 year and 1.2 m in 2 years; growth is stragglier by the 3rd season, when under similar growth conditions plants are thick and bushy, up to 1.5 m high. Growth is poor under natural conditions, 5-8 cm high after 1st season and 17-35 cm after 2nd season; *Z. mauritiana* coppices well and grows vigorously from stumps and root suckers. Fruiting starts after 3-5 years and is usually very abundant.

GERMPLASM MANAGEMENT

Orthodox storage behaviour, viability maintained for 2 years in hermetic air-dry storage at 5 deg. C. The germination rate increases during the 1st year of storage. The cleaned stones can be kept for 5 years in sealed containers, although during this period the viability drops from 95% to 30%. *Z. mauritiana* has 3300 pyrenes/kg.

PESTS AND DISEASES

Fruit flies are a major cause of crop losses, the insects unfortunately having a preference for the same cultivars as humans. Damage by fruit-borers, leaf-eating caterpillars, weevils, leafhoppers and mealy bugs has also been reported. Insect pests include *Meridarchis scyroides*, *Oocussida cruenta*, *Myllocerus* spp., *Thiacidas postica*, *Drepanococcus chiton*, *Florithrips tregardhi* and *Systasis* spp. Powdery mildew can be so serious that leaves and fruitlets drop, but it can be adequately controlled. Lesser diseases are sooty mould, brown rot and leaf-spot.

FURTHER READNG

- Anon. 1986. The useful plants of India. Publications & Information Directorate, CSIR, New Delhi, India.
- Beentje HJ. 1994. Kenya trees, shrubs and lianas. National Museums of Kenya.
- Bekele-Tesemma A, Birnie A, Tengnas B. 1993. Useful trees and shrubs for Ethiopia. Regional Soil Conservation Unit (RSCU), Swedish International Development Authority (SIDA).
- Birnie A. 1997. What tree is that? A beginner's guide to 40 trees in Kenya. Jacaranda designs Ltd.
- Booth FEM, Wickens GE. 1988. Non-timber uses of selected arid zone trees and shrubs in Africa. FAO Conservation Guide. No. 19. Rome.
- Coates-Palgrave K. 1988. Trees of southern Africa. C.S. Struik Publishers Cape Town.
- Dale IR, Greenway PJ. 1961. Kenya trees and shrubs. Buchanan's Kenya Estates Ltd.
- Eggeling. 1940. Indigenous trees of Uganda. Govt. of Uganda.
- Hocking D. 1993. Trees for Drylands. Oxford & IBH Publishing Co. New Delhi.
- Hong TD, Linington S, Ellis RH. 1996. Seed storage behaviour: a compendium. Handbooks for Genebanks: No. 4. IPGRI.
- ICRAF. 1992. A selection of useful trees and shrubs for Kenya: Notes on their identification, propagation and management for use by farming and pastoral communities. ICRAF.
- Jackson JK. 1987. Manual of afforestation in Nepal. Department of Forestry, Kathmandu.
- Jothi BD, Tandon PL. 1995. Present status of insect pests of ber in Karnataka current research. University of Agricultural Sciences, (Banglore). 24:153-155.
- Kokwaro JO. 1976. Medicinal plants of East Africa. East African Literature Bureau.
- Little EL, Wadsworth FH. 1964. Common trees of Puerto Rico and the Virgin Islands. Agricultural Handbook. No. 249. US Department of Agriculture. Washington DC.
- Mbuya LP et al. 1994. Useful trees and shrubs for Tanzania: Identification, Propagation and Management for Agricultural and Pastoral Communities. Regional Soil Conservation Unit (RSCU), Swedish International Development Authority (SIDA).
- Noad T, Birnie A. 1989. Trees of Kenya. General Printers, Nairobi.
- Parkash R, Hocking D. 1986. Some favourite trees for fuel and fodder. Society for promotion of wastelands development, New Delhi, India.
- Perry LM. 1980. Medicinal plants of East and South East Asia : attributed properties and uses. MIT Press. South East Asia.
- Peter G von Carlowitz. 1991. Multipurpose Trees and Shrubs-Sources of Seeds and Inoculants. ICRAF. Nairobi, Kenya
- Schomburg A, Mhango J, Akinnifesi FK. 2001. Marketing of masuku uapaca kirkiana and masawo Ziziphus mauritiana fruits and their potential for processing by rural communities in southern Malawi: Proceeding of the 14th southern Africa regional review and planning workshop, 3-7th September 2001, Harare, Zimbabwe. p.169-176.
- Singh RV. 1982. Fodder trees of India. Oxford & IBH Co. New Delhi, India.
- Sosef MSM, Hong LT, Prawirohatmodjo S. (eds.). 1998. PROSEA 5(3) Timber trees: lesser known species. Backhuys Publishers, Leiden.
- Tembo L, Chiteka ZA, Kadzere I, Akinnifesi FK, Tagwira F. 2008. Blanching and drying period affect moisture loss and vitamin C content in Ziziphus mauritiana Lamk.: African Journal of Biotechnology. 7(8):3100-3106.
- Tembo L, Chiteka ZA, Kadzere I, Akinnifesi FK, Tagwira F. 2008. Storage temperature affects fruit quality attributes of Ber Ziziphus mauritiana Lamk. in Zimbabwe: African Journal of Biotechnology. 7(8):3092-3099.
- Verheij EWM, Coronel RE (eds.). 1991. Plant Resources of South East Asia No 2. Edible fruits and nuts. Backhuys Publishers, Leiden.
- Vimal OP, Tyagi PD. Fuelwood from wastelands. Yatan Publications, New Delhi, India.
- Vogt K. 1995. A field guide to the identification, propagation and uses of common trees and shrubs of dryland Sudan. SOS Sahel International (UK).

Ziziphus mauritiana

Lam.

Rhamnaceae

jujube

Wadsworth FH. 1997. Forest production for tropical America. Agricultural Handbook 710. United States Department of Agriculture.

Williams R.O & OBE. 1949. The useful and ornamental plants in Zanzibar and Pemba. Zanzibar Protectorate.

SUGGESTED CITATION

Orwa C, Mutua A , Kindt R , Jamnadass R, Simons A. 2009. Agroforestry Database:a tree reference and selection guide version 4.0 (<http://www.worldagroforestry.org/af/treedb/>)